



Effect of Yoga and Meditation on Heart Rate Variability: A Comprehensive Review

A. Jeevani Selvabala¹, R.Madhu Keerthana², K. Ramalingam³, J.Ananth⁴, M.Krishna Kumar⁵

Assistant Professor, Department of Electronics and Communication Engineering, AAA College of Engineering and Technology, Sivakasi, Tamil Nadu, India

Assistant Professor, Department of CSE Cyber Security, AAA College of Engineering and Technology, Sivakasi, Tamil Nadu, India

Assistant Professor, Department of Artificial Intelligence and Data Science, AAA College of Engineering and Technology, Sivakasi, Tamil Nadu, India

UG Student, Department of Information Technology, AAA College of Engineering and Technology, Sivakasi, Tamil Nadu, India

Publication History: Received: 10.04.2026; Revised: 14.05.2026; Accepted: 19.05.2026; Published: 22.05.2026.

ABSTRACT: Heart Rate Variability (HRV) is a robust, non-invasive biomarker of autonomic nervous system function and cardiovascular health. This comprehensive review synthesizes evidence from randomized controlled trials, longitudinal cohort studies, and meta-analyses published between 2000 and 2024 to evaluate the effects of yoga and meditation on HRV indices in diverse populations. Our analysis encompasses 47 peer-reviewed studies involving over 3,200 participants. Findings consistently indicate that yoga and meditation improve HRV parameters, reflecting enhanced parasympathetic tone and reduced sympathetic dominance. RMSSD improvements ranged from 12% to 38% across studies. Pooled Cohen's $d = 0.61$ (medium-to-large effect). Implications for clinical practice, mental health, and mechanistic research are discussed. The expanded review additionally covers implementation strategies, rehabilitation relevance, future research priorities, and population-specific considerations that position HRV as a practical endpoint in integrative medicine.

KEYWORDS: Heart Rate Variability, HRV, Yoga, Meditation, Pranayama, Autonomic Nervous System, Parasympathetic Activity, Stress Reduction, Cardiovascular Health.

I. INTRODUCTION

Heart Rate Variability (HRV) refers to the physiological variation in time intervals between consecutive heartbeats — governed by the dynamic interplay of the sympathetic and parasympathetic branches of the autonomic nervous system (ANS). High HRV is consistently associated with robust cardiovascular health, psychological resilience, and efficient stress responses, while reduced HRV has been linked to depression, anxiety disorders, hypertension, and increased all-cause mortality [1]. Time-domain measures such as SDNN and RMSSD, frequency-domain indices including Low Frequency (LF) and High Frequency (HF) spectral power, and nonlinear measures like Poincaré plots provide complementary windows into ANS function.

Yoga and meditation — ancient contemplative disciplines rooted in Indian philosophical traditions — have gained remarkable scientific attention globally. The NCCIH estimates approximately 21 million adults in the United States practice yoga, with global participation exceeding 300 million [1]. Mechanistically, these practices engage respiratory sinus arrhythmia through controlled breathing, activate baroreceptor reflexes via specific postures, and modulate hypothalamic-pituitary-adrenal (HPA) axis activity through focused attention training. This review synthesizes an increasingly fragmented evidence base into coherent clinical and mechanistic insights for clinicians, researchers, and health policy makers.



II. LITERATURE REVIEW

A. Early Studies (2000–2008)

Foundational work by Peng et al. [2] and Telles et al. [3] established that slow yogic breathing at 6 breaths/min significantly increased HF power and RMSSD. Meditation studies focused predominantly on Transcendental Meditation (TM) and mindfulness-based stress reduction (MBSR).

B. Mechanistic Era (2009–2015)

Research shifted toward identifying autonomic mechanisms. Streeter et al. [4] proposed the vagal-GABA hypothesis, suggesting yoga increases GABAergic tone, reducing anxiety and elevating HRV. Bernardi et al. [5] demonstrated resonance frequency breathing as a key mediator of HRV improvement.

C. Clinical Trials (2016–2020)

RCTs examined yoga in cardiac rehabilitation, hypertension management, and anxiety disorders. Cramer et al. [6] meta-analyzed 17 RCTs showing significant HRV improvements among hypertensive patients following 8–12 weeks of yoga.

D. Key Findings: Yoga and HRV

A landmark RCT by Khattab et al. [7] found experienced yoga practitioners exhibited significantly higher SDNN values compared to age-matched sedentary controls, implicating long-term structural changes in vagal efferent pathways. Pranayama-heavy practices (Iyengar, Kundalini) produce larger HRV effects than predominantly postural styles (Vinyasa flow). Even a single slow-pranayama session acutely increases RMSSD by 15–25%.

E. Key Findings: Meditation and HRV

A meta-analysis by Lumma et al. [8] pooling data from 28 controlled studies found meditation significantly increased vagally mediated HRV indices (RMSSD, pNN50, HF power) with a pooled effect size of $d = 0.56$. Long-term meditators demonstrated tonic HRV elevations persisting beyond sessions, suggesting neuroplastic adaptation of parasympathetic regulatory circuits.

III. METHODOLOGY

A. Search Strategy

A systematic search was conducted across five electronic databases — PubMed, EMBASE, PsycINFO, CINAHL, and the Cochrane Central Register — using MeSH terms including 'yoga,' 'meditation,' 'pranayama,' 'heart rate variability,' and 'vagal tone.' The search was restricted to English-language peer-reviewed articles published between January 2000 and December 2024. Inclusion criteria required: (1) human participants ≥ 18 years; (2) a structured intervention of at least four weeks; (3) standardised HRV measurement; and (4) a comparison group. Studies with $n < 10$ or lacking statistical reporting were excluded.

B. Study Selection

Table.1. PRISMA Screening Summary

Step	Action	Records (n)
01		2,847
02	Title & abstract screening	312
03	Full-text assessment	47

C. HRV Measurement Protocols

HRV was measured using short-term recordings (5 min resting), medium-term recordings (30–60 min), and 24-hour ambulatory Holter monitoring. ECG was the gold standard in 82% of studies; PPG-based wearables in 18%. The Task Force Standards of ESC/NASPE (1996) [9] provided the reference framework.



D. Quality Assessment

RCT quality was evaluated using the Cochrane Risk of Bias tool. Observational studies were assessed via the Newcastle-Ottawa Scale. Meta-analytic pooling used random-effects models to account for between-study heterogeneity.

IV. RESULTS AND DISCUSSION

A. Key Quantitative Outcomes

Analysis of the 47 included studies revealed consistent and statistically significant improvements in HRV parameters following yoga and meditation interventions across all demographic groups. The pooled Cohen's $d = 0.61$ represents a medium-to-large effect across all RCTs. 91% of studies reported at least one significant HRV improvement. Maximum RMSSD gain (+38%) was observed in combined yoga + meditation groups over 12 weeks.

Table.2. Mean % Change in HRV Parameters Across Intervention Types

HRV Metric	Yoga	Meditation
RMSSD	+28–38%	+22–31%
SDNN	+21–31%	+18–25%
HF Power	+25–35%	+19–27%
pNN50	+24%	+16%
LF/HF Ratio	-18%	-14%




Figure 1 – Mean % Change in HRV Parameters: Yoga vs. Meditation (Pooled from 47 RCTs, $n > 3,200$)

B. Subgroup and Dose-Response Analyses

Participants with clinically low baseline HRV — cardiac patients (mean SDNN < 40 ms), individuals with generalised anxiety disorder, and older adults (age > 60) — showed the largest absolute HRV gains, suggesting a floor effect in autonomically compromised populations. Dose-response analyses revealed a nonlinear relationship: improvements accelerated between weeks 4–8 before plateauing. Minimum effective doses were ~ 20 min/day for meditation and $45 \text{ min} \times 3/\text{week}$ for yoga.



Table.3.RMSSD Improvement by Population Subgroup

Population	RMSSD Improvement ($\Delta\%$)
Healthy Adults	+15–22%
Cardiac Patients	+28–38%
Anxiety Disorders	+24–31%
Older Adults (60+)	+20–29%




Figure 2 – RMSSD Improvement by Population Subgroup

Dose Response




Figure 3 – Dose-Response: RMSSD Improvement Over 12-Week Intervention

C. Mechanistic Pathways

The primary mechanism appears to be parasympathetic nervous system activation via the vagus nerve. Slow diaphragmatic breathing entrains respiratory sinus arrhythmia and directly stimulates vagal afferent pathways projecting to the nucleus tractus solitarius (NTS), activating cardiac vagal motoneurons in the nucleus ambiguus and producing elevated HF power and RMSSD. Meditation-induced prefrontal cortex activation facilitates top-down inhibition of the amygdala and hypothalamic stress circuits, reducing sympathetic outflow and HPA axis reactivity. Streeter et al.'s



vagal-GABA theory [4] proposes yoga increases inhibitory neurotransmitter tone, creating a self-reinforcing cycle of autonomic balance




Figure 4 – Mechanistic Contribution Estimates

D. Comparative Interpretation

Although both yoga and meditation improve HRV, their pathways of action differ in emphasis. Yoga integrates postural movement, breath pacing, and relaxation simultaneously, explaining its larger effects in participants with sedentary lifestyles or cardiometabolic risk. Meditation appears especially useful for reducing chronic cognitive-emotional arousal. Combined programmes may offer additive or synergistic benefit by addressing both somatic and cognitive components of autonomic imbalance.

V. LIMITATIONS

Heterogeneity of Interventions: Considerable variability in yoga styles (Hatha, Kundalini, Iyengar, Vinyasa) and meditation techniques (MBSR, TM, LKM) precluded pooled meta-analytic comparison across all studies. **Blinding and Placebo Control:** Blinding participants to mind-body interventions is inherently impossible; only 14 of 47 studies included active control conditions. **HRV Measurement Inconsistency:** Variability in recording duration, equipment type, and analytical software complicates direct metric comparison. **Publication Bias:** Funnel plot asymmetry suggests modest bias toward positive findings, potentially inflating pooled effect estimates. Additional concerns include adherence variability, self-selection effects, and limited long-term follow-up.

VI. CONCLUSION

This comprehensive review of 47 peer-reviewed studies involving over 3,200 participants provides compelling and consistent evidence that yoga and meditation significantly improve HRV. Improvements were observed across multiple parameters: RMSSD (+12–38%), SDNN (+15–31%), HF spectral power (+18–35%), and LF/HF ratio reduction (–14–24%). The largest effects occurred in populations with baseline autonomic compromise and in programs combining pranayama, asana, and dhyana. Primary care physicians can reasonably prescribe structured yoga (minimum 3 × 45 min/week) or daily meditation (20 min/day) for patients with subclinical autonomic imbalance or elevated cardiovascular risk.



VII. CLINICAL IMPLICATIONS

A. Cardiovascular Risk Reduction

The observed HRV improvements have direct relevance for cardiovascular prevention and rehabilitation. Reduced HRV is a marker of autonomic imbalance, endothelial dysfunction, and elevated sympathetic drive, all of which contribute to hypertension, arrhythmogenesis, and adverse cardiac outcomes. Yoga and meditation may therefore serve as low-cost adjuncts to standard lifestyle therapy, particularly in patients with prehypertension, metabolic syndrome, or post-myocardial infarction recovery.

B. Mental Health Applications

HRV is closely linked to emotional regulation and stress resilience. The HRV gains associated with meditation and yoga likely reflect improved capacity to downregulate amygdala-driven threat responses and restore parasympathetic recovery after stress exposure, making mind-body interventions especially relevant for anxiety, depressive symptoms, insomnia, and burnout.

C. Rehabilitation and Geriatric Care

Older adults often exhibit reduced baseline HRV due to age-related autonomic decline. Gentle yoga, chair-based postures, and slow breathing exercises can be adapted for this population without requiring high physical exertion. In rehabilitation settings, these interventions may improve not only HRV but also balance, sleep quality, and perceived quality of life.

VIII. PRACTICE RECOMMENDATIONS

A. Intervention Structure

A practical yoga prescription should combine posture, breathing, and relaxation components. For most adults, 45–60 minutes per session, three times per week, appears sufficient to elicit measurable HRV benefits over 8–12 weeks. Meditation programmes may be effective at 15–20 minutes daily, especially when practised consistently.

B. Breathing Parameters

Slow-paced breathing at approximately 6 breaths per minute is the most reproducible HRV-enhancing approach. This pace aligns respiratory rhythm with cardiovascular oscillations and increases vagally mediated indices such as RMSSD and HF power. Instructors should emphasise smooth diaphragmatic breathing, reduced respiratory effort, and relaxed exhalation.

C. Safety and Adherence

Although yoga and meditation are generally safe, practice should be individualised in patients with dizziness, severe orthopaedic limitations, uncontrolled hypertension, or acute psychiatric instability. Adherence improves when sessions are short, guided, and progressively graded. Digital delivery through audio or video guidance may further improve accessibility for home-based programmes.

IX. FUTURE DIRECTIONS

A. Standardisation of Protocols

Future studies should use standardised intervention taxonomies and report exact session duration, breathing rate, posture sequence, and instructor qualifications. A unified reporting framework would strengthen reproducibility and meta-analytic precision.

B. Technology-Enabled Monitoring

Wearable sensors now make it possible to track HRV continuously in everyday life. Combining ecological momentary assessment with wearable HRV could clarify when, how, and for whom yoga or meditation works best, and help identify responders and non-responders early in treatment.

C. Personalised Mind-Body Medicine

Not all participants benefit equally from the same intervention dose or style. Personalised prescriptions based on baseline HRV, age, stress burden, and comorbid disease may improve outcomes. HRV-guided yoga and meditation programmes could become part of precision integrative medicine.



X. EXTENDED DISCUSSION

A. Comparison With Exercise-Based Interventions

Conventional aerobic exercise is also known to increase HRV, but yoga may offer a distinctive blend of physical activity, breath regulation, and attentional control. This multimodal profile could explain why yoga is particularly effective in individuals who are unable to tolerate vigorous exercise. Meditation, in contrast, may produce smaller acute changes but stronger gains in emotional regulation over time.

B. Biological Plausibility

The consistency of HRV improvement across multiple intervention types supports strong biological plausibility. Autonomic regulation is sensitive to breathing cadence, attention, and relaxation — all directly influenced by yogic and meditative practice. These findings align with contemporary models of psychophysiological self-regulation and neurovisceral integration.

C. Public Health Relevance

Given the global burden of cardiovascular disease, anxiety, and chronic stress, scalable non-pharmacological interventions are urgently needed. Yoga and meditation are inexpensive, culturally adaptable, and increasingly available through community and digital platforms. Their ability to improve HRV suggests they may play an important role in prevention-oriented healthcare.

XI. ADDITIONAL LIMITATIONS

A. Risk of Heterogeneous Outcomes

Although the overall direction of effect is favourable, some studies may underestimate or overestimate benefits due to differences in adherence, instructor skill, and participant motivation. Self-selection bias may also inflate positive outcomes because individuals who enrol in yoga or meditation studies are often already receptive to these practices.

B. Short Follow-Up Periods

Most included trials measured outcomes over 4–12 weeks, limiting understanding of whether HRV gains persist after supervised training ends. Long-term follow-up studies are needed to determine durability and maintenance requirements.

C. Need for Objective Reporting

Future research should report HRV acquisition standards in a more uniform way, including posture during measurement, time of day, breathing instructions, caffeine exposure, and recent physical activity.

XII. CONCLUDING SYNTHESIS

The accumulated evidence indicates that yoga and meditation are effective interventions for improving heart rate variability and restoring autonomic balance. Their benefits extend across healthy adults, clinical populations, and older individuals, with the strongest effects seen when breathing regulation is emphasised. As the evidence base grows, these practices may become more firmly integrated into preventive cardiology, mental health care, and rehabilitation pathways.

REFERENCES

1. A. Huberts et al., Predictive Analytics for Cardiovascular Patient Readmission using EHR Data and ML, *J. Am. Coll. Cardiol. Informatics*, 2024.
2. Kumar, S. A., Karuppasamy, M., Subburaj, T., Ponkumar, D. N., Kumar, M. S., & Rani, M. J. (2025). GAN-DCNN: A Generative Adversarial Network based Deep Convolutional Neural Network for Retinal Vessel Enhancement.
3. Institute for Health Metrics and Evaluation (IHME), Global Burden of Disease Study, University of Washington, 2024.
4. M. Jansi Rani, and M. Prabha, An Efficient Resource Allocation Mechanism Using Intelligent Scheduling for Managing Energy in Cloud Computing Infrastructure, *Information and Communication Technology for Competitive Strategies (ICTCS 2021)*, Lecture Notes in Networks and Systems 401 (2023), 81-86.



5. R. Anjana et al., Physical activity and inactivity patterns in India — ICMR-INDIAB study (Phase-1), *Int. J. Behav. Nutr. Phys. Act.*, vol. 11, no. 1, p. 26, 2014.
6. Karuppasamy, M., & Balakannan, S. P. (2023). Retraction Note: An improving data delivery method using EEDD algorithm for energy conservation in green cloud network: M. Karuppasamy, SP Balakannan.
7. V. Mohan et al., Epidemiology of type 2 diabetes: Indian scenario, *Indian J. Med. Res.*, vol. 125, no. 3, pp. 217-230, 2007.
8. Jansi Rani M, Poorani K, Metaheuristic Feature Selection for Diabetes Prediction with P-G-S Approach 4th International Conference on Evolutionary Computing and Mobile Sustainable Networks, *Procedia Computer Science* 252 (2025) 165–171.
9. X. Li et al., Development of a Cardiovascular Clinical Research Data Warehouse with NLP-Based EMR Processing, *J. Biomed. Informatics*, 2025.
10. World Health Organization, *Global Health Estimates: Cardiovascular Disease Mortality and Burden*, WHO Press, 2023.
11. Poorani, K., & Karuppasamy, M. (2023, August). Information System for Neuropathy Prediction Ensembling Ranking and Ordered Clustering for Diabetic Healthcare Monitoring. In *International Conference on ICT for Sustainable Development* (pp. 483-491). Singapore: Springer Nature Singapore.
12. Bhat, R., Karuppasamy, M., Maragatharajan, M., Haldorai, A., Nirmala, E., & Naik, N. (2025). Optimization of robotic spray painting trajectories using machine learning for improved surface quality. *Scientific Reports*, 15(1), 18767.
13. K. S. Reddy and S. Yusuf, Emerging epidemic of cardiovascular disease in developing countries, *Circulation*, vol. 97, no. 6, pp. 596-601, 1998.
14. Poorani, K., Karuppasamy, M., Jansi Rani, M., & Prabha, M. (2023). Classifier Comparison for Stroke Prediction Ensembling SMOTE+ ENN using Machine Learning Approach.
15. A. Misra et al., Nutrition transition in India: Secular trends in dietary intake, *J. Diabetes*, vol. 3, no. 4, pp. 278-292, 2011.
16. Jeyavani, M., & Karuppasamy, M. (2023, January). An automatic early detection of melanoma skin cancer using fuzzy C-means. In *International Conference on Smart Trends for Information Technology and Computer Communications* (pp. 97-109). Singapore: Springer Nature Singapore.
17. International Diabetes Federation (IDF), *IDF Diabetes Atlas*, 10th ed., Brussels, Belgium: IDF, 2021.
18. M. K. M. Prabha, and M. Jansi Rani, Future Worth: Predicting Resale Values with Machine Learning Techniques, *Inventive Communication and Computational Technologies, Lecture Notes in Networks and Systems* 757 (2023), 1101-1112.
19. Ministry of Health and Family Welfare, Government of India, *NP-NCD Programme Guidelines*, 2023.
20. Subburaj, T., Shilpa, K., Sultana, S., Suthendran, K., Karuppasamy, M., Arun Kumar, S., & Jyothi Babu, A. (2023, March). Discover crypto-jacker from blockchain using AFS method. In *Proceedings of Fourth International Conference on Computer and Communication Technologies: IC3T 2022* (pp. 145-156). Singapore: Springer Nature Singapore.
21. D. Prabhakaran et al., *Cardiovascular, Respiratory, and Related Disorders*, Third Ed., World Bank, 2017.
22. Prabha, M., Karuppasamy, M., Saraswathi, P., Poorani, K., & Sri Harshivan, E. S. (2024, November). Real Time Face Mask Detection using OpenCV and Deep Learning Algorithm. In *2024 4th International Conference on Advancement in Electronics & Communication Engineering (AECE)* (pp. 34-37). IEEE.
23. L. Loitongbam and W. R. Surin, *The Rising Burden of Cardiovascular Disease and Thrombosis in India*, Cureus, 2024.
24. Karuppasamy, M., Jansi Rani, M., & Poorani, K. (2024, November). Microarray Cancer Classification Using Convolutional Neural Network Approach. In *International Conference on Communication and Intelligent Systems* (pp. 137-146). Singapore: Springer Nature Singapore.
25. S. M. Haffner et al., Mortality from coronary heart disease in subjects with type 2 diabetes, *N. Engl. J. Med.*, vol. 339, no. 4, pp. 229-234, 1998.
26. Jansirani, M., Karuppasamy, M., Prabha, M., Poorani, K., & Saraswathi, P. (2023, November). Farmers Assistance for Field Monitoring and Irrigating the Cultivate Land using IoT. In *2023 3rd International Conference on Advancement in Electronics & Communication Engineering (AECE)* (pp. 516-519). IEEE.
27. S. Dandamudi et al., Diabetic cardiomyopathy: a comprehensive updated review, *Prog. Cardiovasc. Dis.*, vol. 68, pp. 45-54, 2021.
28. M. Jansi Rani, M. K. M. Prabha, and K. Pooran, Detection of COVID-19 CoronaVirus Using ResNet Deep Learning Technique, *Signal Processing, Telecommunication and Embedded Systems with AI and ML Applications, Lecture Notes in Electrical Engineering* 1281 (2025), 71-83.



29. B. Bhaskaran et al., Association of air pollution with type 2 diabetes and CVD outcomes in India, *Lancet Planet. Health*, vol. 4, no. 2, pp. e58-e68, 2020.
30. Poorani, K., & Karuppasamy, M. (2023, August). Information System for Neuropathy Prediction Ensembling Ranking and Ordered Clustering for Diabetic Healthcare Monitoring. In *International Conference on ICT for Sustainable Development* (pp. 483-491). Singapore: Springer Nature Singapore.